

THE SIGNATURE OF A MANIFOLD

José Rodríguez

Abstract

Let us consider a compact oriented riemannian manifold M without boundary and of dimension $n = 4k$. The signature of M is defined as the signature of a given quadratic form Q . Two different products could be used to define Q and they render equivalent definitions: the exterior product of forms and the cup product of cohomology classes. The signature of a manifold is proved to yield a topological invariant. Additionally, using the metric, a suitable Dirac operator can be defined whose index coincides with the signature of the manifold.

1 Introduction

Let us imagine a rubber surface filled in electric charges with a given density. Clearly, if we deform the surface, the total electric charge is conserved. Thus, thanks to electric charges we have an invariant. The question is whether or not we can somehow balance the role of the densities of charge in such a way that we get a charge-free invariant, i.e., an invariant that depends only on the manifold. In the following we present a positive partial answer to this question.

In differential geometry we can integrate forms and only forms. The dimension of the form is given by the dimension of the domain of integration. Say, if we have a three dimensional density of electric charge, the total charge is its integral: the charge density shall be encoded as a 3-form. Likewise, if we want to reproduce the fact that the line integral of a force gives a work, we encode forces as 1-forms. On the other hand, Force = Electric field $\times$ charge, so that the electric field could be encoded as a 1-form also. The Lorentz force is a 1-form given by $\vec{F} = k\vec{E} + c\vec{B} \times \vec{v}$, therefore $\vec{B}$ is encoded as a 2-form that operates over a pair of vectors, $\vec{v}$ and a displacement.

We know that the divergence of a rotational is zero, and that the rotational of a gradient is also zero. These facts lead to the general law $d^2 = 0$ which more exactly means: Let d_k denote the exterior derivative of k -forms, then $d_{k+1} \circ d_k = 0$.

The global object to be integrated is a form which is an assignation of a k -differential form to everyone point of the manifold. The space of forms is denoted as $\Omega(M)$. So, a charge density is a 3-form, i.e., an element of $C^\infty(M, \Lambda^3 T^*(M)) = \Omega^3(M)$. We use standard notation $[N], [F]$.

Stokes' theorem over a manifold M reads: $\int_{\partial M} \omega = \int_M d\omega$. This theorem is a generalization of Gauss and Stokes theorems of vector calculus: what ω causes

on ∂M , the border of M , is completely evidenced by studying what $d\omega$ causes in the interior of M .

Definitions. Let d_k denote the exterior derivative of k -forms and δ_{k+1} its adjoint. We have that $d_{k+1} \circ d_k = 0$. The subindex of the exterior derivative rarely is explicitly written and d_k is simplified into d and $d_{k+1} \circ d_k = 0$ is noted simply as $d^2 = 0$. The same abuse of notation will be applied to other operators that appear in the sequel. A form ω is called closed if $d\omega = 0$, coclosed if $\delta\omega = 0$, exact if there exist ψ such that $d\psi = \omega$, coexact if there exists ϕ such that $\delta\phi = \omega$.

Since $d_{k+1} \circ d_k = 0$ then $Im(d_k) \subseteq Ker(d_{k+1})$ and moreover both are groups. The quotient $H^k = Ker(d_{k+1})/Im(d_k)$ is a group that is called the de Rham k -cohomology group.

If ω_1 and ω'_1 belong to the same cohomology class then they belong in the kernel of d_{k+1} , therefore $d_{k+1}\omega = d_{k+1}\omega' = 0$ and moreover they differ by an exact form $d_k\tau$, so $\omega_1 = \omega'_1 + d_k\tau$ with $d_{k+1} \circ d_k\tau = 0$. The operator d is antisymmetric. Hence, $d_k\omega = 0$ for $k > n$ given that n is the dimension of the manifold, $H^k(M) = 0$ for $k > n$.

In the light of the Poincaré's lemma, the de Rham cohomology group is regarded as an obstruction to the global exactness of closed forms. The dimension of a cohomology group is finite and is known as the Betty number of the group and therefore could be seen as a measure of the variability of the global inexactness of closed forms. The existence of electric motors is due to the fact that $H^1(\mathbb{R}^n - \{0\})$ is not trivial. Similar effects are observed in the theory of fundamental interactions.

2 The signature quadratic form

Let M be a compact oriented riemannian manifold without boundary and of dimension $n=4k$. The signature quadratic form of M is defined to be the bilinear form

$$Q : H^{2k}(M) \otimes H^{2k}(M) \rightarrow \mathbb{R}$$

$$Q([\omega_1], [\omega_2]) = \int \omega_1 \wedge \omega_2$$

The signature quadratic form is well defined, i.e., the integral on the right does not depend on the representatives of the cohomology classes. To see this, let us take another representative of $[\omega_1]$ say ω'_1 . Now, $d\omega_1 = d\omega'_1 = d\omega_2 = 0$ and ω_1 and ω'_1 differ by an exact form $d\tau$, where τ is a $(2k-1)$ -form. So, $\omega_1 = \omega'_1 + d\tau$ and $d^2\tau = 0$.

Then

$$\int \omega'_1 \wedge \omega_2 = \int (\omega_1 + d\tau) \wedge \omega_2 = \int \omega_1 \wedge \omega_2 + \int d\tau \wedge \omega_2$$

Recalling that $d\omega_2 = 0 = \tau \wedge d\omega_2$ and using Leibniz's rule and Stokes' theorem over a manifold without boundary we have:

$$\int d\tau \wedge \omega_2 = \int_M d\tau \wedge \omega_2 = \int_M d\tau \wedge \omega_2 \pm \int_M \tau \wedge d\omega_2 = \int_M d(\tau \wedge \omega_2) = \int_{\partial M} \tau \wedge \omega_2 = 0$$

and henceforth

$$\int \omega'_1 \wedge \omega_2 = \int \omega_1 \wedge \omega_2.$$

Extending the reasoning to the general form of representatives of $[\omega_2]$ we obtain the independence of the bilinear form from representatives and so Q is defined over cohomology classes of order $2k$.

Theorem. The signature quadratic form Q is symmetric iff $n=4k$.

Proof: In general we have:

$\omega \wedge \nu = (-1)^{pq} \nu \wedge \omega$ when ω is a p -form and ν is a q -form. Thus, if $p = q$ then $(-1)^{pq} = (-1)^{p^2}$. But $p^2 = p \bmod 2$, i.e., $p^2 - p = p(p-1) = 2m$, i.e., either p or $p-1$ is even and hence $p(p-1)$ is always even. So, $(-1)^{p^2} = (-1)^p = 1$ iff p is even iff $= 2p$ is multiple of 4.

2.1 The cup product

Since the wedge product is defined for pairs of forms of any order we can verify that the vector space $H^* = H^*(M) = \bigoplus_{p \geq 0}^n H^p M$ can be endowed with a ring structure as follows: for $[\omega] \in H^p$ and $[\nu] \in H^q$ we define the cup product as

$$\cup : H^* \times H^* \rightarrow H^*$$

$$[\omega] \cup [\nu] = [\omega \wedge \nu]$$

Let us check that this is a well defined product in the set of cohomology classes: if $[\omega] \in H^p$ and $[\nu] \in H^q$ then $\omega \wedge \nu \in \Omega^{p+q}$ and $[\omega \wedge \nu] \in H^{p+q}$ but we shall show that if we take other representatives their wedge product is still in $[\omega \wedge \nu]$.

If $\omega, \omega' \in [\omega]$ in H^p then $d\omega = d\omega' = 0$ and $\omega' = \omega + d\tau$. If $\nu \in H^q$ then $d\nu = 0$. Therefore $\omega' \wedge \nu = (\omega + d\tau) \wedge \nu = \omega \wedge \nu + d\tau \wedge \nu = \omega \wedge \nu + d(\tau \wedge \nu)$ because $d(\tau \wedge \nu) = d\tau \wedge \nu + \tau \wedge d\nu = d\tau \wedge \nu$. Henceforth, $\omega' \wedge \nu$ and $\omega \wedge \nu$ differ by an element of the form $d(\alpha)$ where $\alpha = \tau \wedge \nu$ and so they belong to the same cohomology class.

Restricting the cup product to $H^{2k}(M) \otimes H^{2k}(M)$, the signature quadratic form Q takes the form:

$$Q([\omega_1], [\omega_2]) = \int [\omega_1] \cup [\omega_2] = \int \omega_1 \wedge \omega_2$$

The integral of any p -form has an intrinsic definition, i.e., it is devised to be invariant under any change of coordinates. Therefore, Q is also intrinsically defined. Hence, to calculate it we can take anyone system of coordinates on M . On the other hand, to say that Q is symmetric is the same as to say that the cup product is commutative when restricted to $H^{2k}(M) \otimes H^{2k}(M)$.

2.2 The signature of a manifold

Since the space H^{2k} is finite dimensional with dimension b_{2k} , called the $2k$ -th Betti number, we can take a basis $E = \{[h_i]\}$ with respect to which the quadratic form Q has an associated matrix, $Q_{[E]}$, whose entries are $Q_{[E],ij} = \int [h_{*i}] \wedge [h_{*j}]$. This matrix can be diagonalized to a matrix with eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_s$, where $s = \dim H^{2k} = b_{2k}$. Below we will see a concrete basis in which Q is diagonal. The signature of $Q_{[E]}$ is the number of positive eigenvalues minus the number of negative eigenvalues. Under a change of basis with the same orientation in H^{2k} we get, up to a permutation, the same eigenvalues. All our results are summarized in the following

Definition. The signature of compact, without boundary, orientable manifold, $\sigma(M)$, is the signature of the quadratic form Q . The signature of the manifold is the number of positive eigenvalues minus the number of negative eigenvalues of the finite dimensional matrix of Q in any basis of H^{2k} and in any system of coordinates.

3 Invariance

When two manifolds could be smoothly deformed back and forth, their signatures coincide. Technically, the signature of a manifold is invariant under diffeomorphisms. This property is expected because a diffeomorphism represents a change of coordinates and we have seen that the signature of a manifold is an intrinsic object so that it is independent of coordinates. A direct demonstration over the ring H^* is instructive and can easily be carried out thanks to the pull-back technology. When in calculus one says "change of variable" in differential geometry one says "pull-back" of p -forms, which is built upon the notion of differential of a function:

Definitions. Let $\phi : M \rightarrow N$ be a map of manifolds and let $\phi(x) = y$. Let $T_x M$ and $T_{\phi(x)} N$ be the tangent spaces at x over M and, respectively, at $\phi(x)$ over N . We define the differential ϕ_* of ϕ as the linear isomorphism $\phi_* : T_x M \rightarrow T_{\phi(x)} N$ such that for any vector $v \in T_x M$ we have $[\phi_*(v)](f) = v(\phi \circ f)$ for any scalar function $f : N \rightarrow \mathbb{R}$. A map is called differentiable or smooth if its differential exists. The pull-back ϕ^* of a smooth map $\phi : M \rightarrow N$ is the linear transformation $\phi^* : \Lambda_{\phi(x)}^1 N \rightarrow \Lambda_x^1 M$ such that $\phi^*(\omega)(v) = \omega(\phi_*(v))$ for all vectors $v \in T_x M$ and 1-forms ω .

The pull back observes the following properties, the first of which allows to extend the pull back of 1-forms to the whole space of form:

- 1) $\phi^*(\alpha \wedge \beta) = \phi^*(\alpha) \wedge \phi^*(\beta)$
- 2) $\phi^*(d\alpha) = d(\phi^*(\alpha))$
- 3) $(\phi^{-1})^* = (\phi^*)^{-1}$

Lema. Let $\phi : M \rightarrow N$ be an orientation preserving diffeomorphism between manifolds and let $H^*(M)$ and $H^*(N)$ the respective cohomology rings, then ϕ

induces a ring contravariant isomorphism $F_\phi^* : H^*(N) \rightarrow H^*(M)$ defined by $F_\phi^*([\eta]) = [\phi^*(\eta)]$.

Proof: Let us prove that F_ϕ^* is well defined, with an inverse which is the ring homomorphism induced by ϕ^{-1} and that $F_\phi^*([\eta_1] \cup [\eta_2]) = F_\phi^*([\eta_1]) \cup F_\phi^*([\eta_2])$.

To see that F_ϕ^* is well defined we need to prove that two members of the same cohomology class in the domain are transformed into members of the same cohomology class in the range. Two elements are in the same cohomology class if they differ by an exact form, i.e., if $[\eta_1] = [\eta_2]$ then $\eta_1 = \eta_2 + d\tau$. In this case,

$$\phi^*(\eta_1) = \phi^*(\eta_2) + \phi^*(d\tau) = \phi^*(\eta_2) + d(\phi^*(\tau))$$

this means that the members of a class are transformed into elements that differ by an exact form, i.e., they are members of the same class:

if $[\eta_1] = [\eta_2]$ then $[\phi^*(\eta_1)] = [\phi^*(\eta_2)]$ so that $F_\phi^*([\eta_1]) = F_\phi^*([\eta_2])$ is well defined.

Let us verify now that if $F_\phi^*([\eta]) = [\phi^*(\eta)]$, the inverse of F_ϕ^* is the ring homomorphism $G_{\phi^{-1}}^*$ induced by ϕ^{-1} , i.e., if $G_{\phi^{-1}}^*([\omega]) = [(\phi^{-1})^*(\omega)]$ then $G_{\phi^{-1}}^*(F_\phi^*([\eta])) = [\eta]$ and $F_\phi^*(G_{\phi^{-1}}^*([\omega])) = [\omega]$. For the first part we have:

$$G_{\phi^{-1}}^*(F_\phi^*([\eta])) = G_{\phi^{-1}}^*([\phi^*(\eta)]) = [(\phi^{-1})^*(\phi^*(\eta))] = [(\phi^*)^{-1}(\phi^*(\eta))] = [I\eta] = [\eta].$$

The proof of the second part is similar.

Let us show that F_ϕ^* is indeed a ring homomorphism:

$$\begin{aligned} F_\phi^*([\eta_1] \cup [\eta_2]) &= F_\phi^*([\eta_1 \wedge \eta_2]) = [\phi^*(\eta_1 \wedge \eta_2)] = [\phi^*(\eta_1) \wedge \phi^*(\eta_2)] \\ &= [\phi^*(\eta_1)] \cup [\phi^*(\eta_2)] = F_\phi^*([\eta_1]) \cup F_\phi^*([\eta_2]). \end{aligned}$$

In conclusion, F_ϕ^* is a ring homomorphism:

$$F_\phi^*([\eta_1] \cup [\eta_2]) = F_\phi^*([\eta_1]) \cup F_\phi^*([\eta_2]).$$

We also say that F_ϕ^* is orientation preserving, in the sense that if we choose a basis E of $H^{2k}(N)$ then $F_\phi^*(E)$ will be a basis of $H^{2k}(M)$ that observe the same orientation.

Convention. The ring isomorphism F_ϕ^* is denoted as ϕ^* .

Theorem. The quadratic signature form Q is invariant under diffeomorphisms.

Proof. Let $\phi : M \rightarrow N$ be a diffeomorphism between manifolds, and let F_ϕ^* be its induced isomorphism between $H^*(M)$ and $H^*(N)$. The theorem of change of variables reads:

$$\int_{N=\phi(M)} \eta = \int_M \phi^*(\eta).$$

Let us apply this theorem to the signature quadratic form:

$$\begin{aligned} \int_{N=\phi(M)} [\eta_1] \cup [\eta_2] &= \int_{N=\phi(M)} \eta_1 \wedge \eta_2 = \int_M \phi^*(\eta_1 \wedge \eta_2) = \int_M \phi^*\eta_1 \wedge \phi^*\eta_2 = \\ &= \int_M [\phi^*\eta_1 \wedge \phi^*\eta_2] = \int_M [\phi^*\eta_1] \cup [\phi^*\eta_2] = \\ &= \int_M F_\phi^*([\eta_1]) \cup F_\phi^*([\eta_2]). \end{aligned}$$

Thus, we have proved that $\int_{N=\phi(M)} [\eta_1] \cup [\eta_2] = \int_M F_\phi^*([\eta_1]) \cup F_\phi^*([\eta_2])$ and this means that two diffeomorphic manifolds have the same quadratic signature form:

$$Q([\phi^*\omega_1], [\phi^*\omega_2]) = Q([\omega_1], [\omega_2]).$$

Corollary. Diffeomorphic oriented manifolds have the same signature.

Proof: Let N be a manifold and $Q([\omega_1], [\omega_2])$ its signature form. The signature $\sigma(N)$ of N is the number of positive eigenvalues minus the number of negative eigenvalues of the finite dimensional matrix of Q in any basis of $H^{2k}(N)$ and in any system of coordinates. Let $\phi : M \rightarrow N$ be a diffeomorphism between manifolds that preserves orientation. The previous theorem shows that they have the same quadratic form $Q([\phi^*\omega_1], [\phi^*\omega_2]) = Q([\omega_1], [\omega_2])$.

Moreover, the preimage of a basis E of $H^{2k}(N)$ through the orientation preserving isomorphism among rings ϕ^* is also a basis with the same orientation. So, we can express $Q([\phi^*\omega_1], [\phi^*\omega_2])$ in the basis $\phi^*(E)$ of $H^{2k}(M)$ and count the difference between positive and negative eigenvalues. We claim that this difference is precisely the signature of M , because we have the freedom of choosing the basis in which Q is expressed.

3.1 Invariance of $\sigma(M)$ under homotopy equivalence

All our study is motivated by the conservation of the total electric charge of a charged body under deformations. If the body is three dimensional and of light metal, without much effort we could smash it to produce a surface. With due care, the total charge would not change. What we are ready to observe is that this case is also admitted in our mathematical formalism.

Definition. Two functions with the same domain and codomain, $h_0, h_1 : Z \rightarrow W$, are homotopically equivalent, $h_0 \sim h_1$, if there exists a smooth map $F : Z \times [0, 1] \rightarrow W$ such that $F|_{Z \times \{0\}} = h_0, F|_{Z \times \{1\}} = h_1$.

The functions shall not be onto and the images of the two functions could be disjoint. Intuitively, two functions are homotopically equivalent if there exists a continuous deformation of $h_0(Z)$ into $h_1(Z)$.

Definition. Let M and N be two oriented smooth compact manifolds. We say that M and N are (strongly) homotopically equivalent if there exist two orientation preserving smooth maps

$$g : N \rightarrow M$$

$$f : M \rightarrow N$$

such that $f \circ g \sim id_N$ and $g \circ f \sim id_M$.

To understand the meaning of this concept let us imagine that M is a person and uses N as a mirror to observe herself. If M can deform herself smoothly into her image $g(f(M))$ and if N also can do the same in her side, then, we say that the two manifolds are homotopically equivalent.

The most important difference between isomorphic manifolds and homotopically equivalent manifolds are considered in the next definitions.

Definition. Let R be a, not empty, subspace of M . If there exists a continuous map $f : M \rightarrow R$ such that $f|_R = id_R$, R is called a retract of M and f a retraction.

Definition. Let R be a, not empty, subspace of M and f a retraction of M over R . R is said to be a deformation retract if id_M and f are homotopically equivalent and R is point by point invariant in the deformation.

The aim of this section is to show that the signature of a manifold is conserved under homotopy deformations including retractions. We shall highlight the fact that our retractions eliminate homotopic redundances but holes are not eliminated. So, a circle cannot be joined to its north pole by a deformation retraction.

To prove that the signature of a manifold is conserved under homotopy equivalence we shall guarantee that homotopy equivalence does not distort cohomology groups. This implies that the antitransport of closed forms through two homotopy equivalent functions belong to the same cohomology class, i.e., that they differ by an exact form.

Lemma. Let $f, g : M \rightarrow N$ be smooth maps that are homotopic to each other. If $\omega \in \Omega^K(N)$ is a closed form, the difference of the pullback images is exact:

$$f^*\omega - g^*\omega = d\psi$$

where $\psi \in \Omega^{k-1}(M)$ and $f^*, g^* : H^*(N) \rightarrow H^*(M)$.

Proof. Since $f \sim g$, there exists a smooth map $F : M \times [0, 1] \rightarrow N$ such that $F|_{M \times \{0\}} = f, F|_{M \times \{1\}} = g$, i.e., $F(x, 0) = f(x)$ and $F(x, 1) = g(x)$ for $x \in M$.

Now we will involved in a game that uses F and the fundamental theorem of calculus: to prove that $f^*\omega - g^*\omega = d\psi$ for some ψ , we will prove a rigorous version of the following idea: $f^*\omega - g^*\omega = d \int F^*\omega$. In what follows, the integral will be replaced by the operator P .

Let us consider a k -form $\eta \in \Omega^k(M \times [0, 1])$. η takes the form $\eta = a_{i_1 \dots i_k}(x, t) dx^{i_1} \wedge \dots \wedge dx^{i_k} + b_{j_1 \dots j_{k-1}}(x, t) dt \wedge dx^{j_1} \wedge \dots \wedge dx^{j_{k-1}}$ where $x \in M, t \in [0, 1]$. The second term is of degree k but it shall include dt so it has only $k - 1$ degrees of freedom to choose its components.

Define a map $P : \Omega^k(M \times I) \rightarrow \Omega^{k-1}(M)$ by

$$P(\eta) = (\int_0^1 b_{j_1 \dots j_{k-1}}(x, s) ds) dx^{j_1} \wedge \dots \wedge dx^{j_{k-1}}$$

Let f_t be a map $f_t : M \rightarrow M \times I$ such that $f_t(p) = (p, t)$

We have

$$f_t^* \eta = a_{i_1 \dots i_k}(x, t) dx^{i_1} \wedge \dots \wedge dx^{i_k} \in \Omega^k(M)$$

since $f_t^*(dt \wedge dx^{j_1} \wedge \dots \wedge dx^{j_{k-1}}) = 0$.

Let us prove now that

$$d(P(\eta)) + P(d\eta) = f_1^*(\eta) - f_0^*\eta.$$

Indeed, if we calculate each term in the lhs we get

$$\begin{aligned} d(P(\eta)) &= d\left(\int_0^1 b_{j_1 \dots j_{k-1}}(x, s) ds\right) dx^{j_1} \wedge \dots \wedge dx^{j_{k-1}} \\ &= \left(\int_0^1 (\partial b_{j_1 \dots j_{k-1}}(x, s) / \partial x^{j_k}) ds\right) dx^{j_k} \wedge dx^{j_1} \wedge \dots \wedge dx^{j_{k-1}} \end{aligned}$$

On the other hand

$$\begin{aligned} P(d\eta) &= P[d(a_{i_1 \dots i_k}(x, t) dx^{i_1} \wedge \dots \wedge dx^{i_k} + b_{j_1 \dots j_{k-1}}(x, t) dt \wedge dx^{j_1} \wedge \dots \wedge dx^{j_{k-1}})] \\ &= P[(\partial a_{i_1 \dots i_k}(x, t) / \partial x^{i_{k+1}}) dx^{i_{k+1}} \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k} \\ &\quad + (\partial a_{i_1 \dots i_k}(x, t) / \partial t) dt \wedge dx^{i_1} \wedge \dots \wedge dx^{i_k}] \\ &\quad + (\partial b_{j_1 \dots j_{k-1}}(x, t) / \partial x^{j_k}) dx^{j_k} \wedge dt \wedge dx^{j_1} \wedge \dots \wedge dx^{j_{k-1}}) \\ &= \left(\int_0^1 (\partial a_{i_1 \dots i_k}(x, s) / \partial s) ds\right) dx^{i_1} \wedge \dots \wedge dx^{i_k} \\ &\quad - \left(\int_0^1 (\partial b_{j_1 \dots j_{k-1}}(x, s) / \partial x^{j_k}) ds\right) dx^{j_k} \wedge dx^{j_1} \wedge \dots \wedge dx^{j_{k-1}} \end{aligned}$$

Summing up these two terms we get

$$d(P(\eta)) + P(d\eta) = \left(\int_0^1 (\partial a_{i_1 \dots i_k}(x, s) / \partial s) ds\right) dx^{i_1} \wedge \dots \wedge dx^{i_k}.$$

Applying the fundamental theorem of calculus for a continuous function we have

$$\begin{aligned} d(P(\eta)) + P(d\eta) &= [a_{i_1 \dots i_k}(x, 1) - a_{i_1 \dots i_k}(x, 0)] dx^{i_1} \wedge \dots \wedge dx^{i_k} \\ d(P(\eta)) + P(d\eta) &= f_1^*(\eta) - f_0^*\eta \end{aligned}$$

Let us apply this identity to the pull back of a closed form $\omega \in \Omega^k(N)$ and $\eta = F^*\omega \in \Omega^k(M \times [0, 1])$:

so $d(P(\eta)) + P(d\eta)$ becomes

$$d(P(F^*\omega)) + P(dF^*\omega) = f_1^*(F^*\omega) - f_0^*(F^*\omega)$$

recalling that $(FG)^* = G^*F^*$ we get

$$d(P(F^*\omega)) + P(dF^*\omega) = (Ff_1)^*\omega - (Ff_0)^*\omega$$

but $f_t(x) = (x, t)$ and moreover $f \sim g$ so that $F(f_0(x)) = F(x, 0) = f(x)$

and $F(f_1(x)) = F(x, 1) = g(x)$

Hence

$$d(P(F^*\omega)) + P(dF^*\omega) = f^*\omega - g^*\omega.$$

Recalling now that ω was chosen to be closed, $d\omega = 0$ and therefore $F^*d\omega = 0$. We can rewrite this as $d(F^*\omega) = 0$ because the pull back and the exterior derivative commute. Integrating with P between 0 and 1 we have $P(dF^*\omega) = 0$.

Replacing

$$d(P(F^*\omega)) + P(dF^*\omega) = d(P(F^*\omega)) = f^*\omega - g^*\omega$$

Explicitly $f^*\omega - g^*\omega = d\psi$ where $\psi = P(F^*\omega)$ proving that the difference of the pullbacks of closed forms throughout homotopy equivalent functions is exact.

Corollary. Let $f, g : M \rightarrow N$ be maps which are homotopic to each other. Then, the pullback maps f^* and g^* defined on the de Rham cohomology groups are identical.

Equivalently, the cohomology class of the difference of the pullback of a closed form through two homotopy equivalent functions is the 0 cohomology class:

$$[f^*\omega - g^*\omega] = [f^*\omega] - [g^*\omega] = [d\psi] = 0$$

Theorem. If M and N are homotopy equivalent even oriented compact manifolds, then $\sigma(M) = \sigma(N)$.

Proof. Since M and N are homotopy equivalent manifolds, there exist two orientation preserving smooth maps

$$f : M \rightarrow N$$

$$g : N \rightarrow M$$

such that $f \circ g \sim id_N$ and $g \circ f \sim id_M$.

It is clear that $g(f(M))$ is a retract of M and that $f(g(N))$ is a retract of N . Let us define the (f,g) -core of M as $Core_{(f,g)}(M) = \bigcap_n (g \circ f)^n(M)$ and the $Core_{(f,g)}(N) = \bigcap_n (f \circ g)^n(N)$. We have at last that the great manifolds can be compressed to their cores and that all of them have isomorphic cohomology rings, which uniquely determines the signature of the manifolds.

Disclaimer. Two manifolds could be inequivalent and yet they can have the same cohomology groups and henceforth the same signature. Example: take as $M = S^2 \times S^4$ and as $N = CP^3$.

Proof. Unknown.

3.2 The Euler characteristic

Let $*$ note the Hodge-star operator, which associates to anyone p -form ω the one and only $(n-p)$ -form $*\omega$ that complements ω to form the volume form: $\omega \wedge (*\omega) = \alpha vol$ where vol denotes de volume form over the manifold and the scalar α is the inner product of the two forms: given two p -forms, ω, η , their inner product is defined as

$$(\omega, \eta) = (1/r!) \int_M \omega_{\mu_1 \dots \mu_r} \eta^{\mu_1 \dots \mu_r} \sqrt{g} dx^1 \dots dx^n = \int \omega \wedge *\eta.$$

Because $** = (-1)^{k(n-k)}$, when operating over k -forms, there is a natural bijection between H^k and H^{n-k} known as Poincaré' duality. Let $b_k = \dim H^k$. Define the Euler characteristic of M as $\chi(M) = \sum (-1)^k b_k$.

Lemma. Let $\dim M = 4k$ then the Euler characteristic of M and the dimension of H^{2k} have the same parity, i.e., $\chi(M) = b_{2k} \bmod 2$.

Proof. $\chi(M) = \sum (-1)^k b_k = \sum_0^{2k-1} (-1)^k b_k + (-1)^{2k} b_{2k} + \sum_{2k+1}^{4k} (-1)^k b_k$.

Using the Poincaré duality, this can be rewritten as

$$\chi(M) = 2 \sum_0^{2k-1} (-1)^k b_k + (-1)^{2k} b_{2k}$$

$$\chi(M) - b_{2k} = 2 \sum_0^{2k-1} (-1)^k b_k$$

that shows that at both sides of this equation we are dealing with even numbers. Or, equivalently, $\chi(M) = b_{2k} \bmod 2$.

3.3 Harmonic forms

We consider here a closed manifold with a metric (M, g) .

Definitions. The Dirac operator D is $D = d + \delta$ and the Laplacian $\Delta = D^2 = (d + \delta)^2 = (d + \delta)(d + \delta) = d^2 + d\delta + \delta d + \delta^2 = d\delta + \delta d$. A form that satisfies the Laplace equation $\Delta\omega = 0$ is called harmonic. The space of harmonic forms of degree k is denoted as $Harm^k(M)$.

Theorem. Δ is, formally, selfadjoint in $L^2(\Lambda T^*M)$.

Proof: $(\Delta\omega, \psi) = ((d\delta + \delta d)\omega, \psi) = (d\delta\omega, \psi) + (\delta d\omega, \psi) = (\delta\omega, \delta\psi) + (d\omega, d\psi) = (\omega, d\delta\psi) + (\omega, \delta d\psi) = (\omega, d\delta\psi + \delta d\psi) = (\omega, (d\delta + \delta d)\psi) = (\omega, \Delta\psi)$.

A full fledged proof requires considerations about the domain of operators [P].

Theorem. $\dim Harm_k(M) < \infty$.

Proof: Theory of Elliptic operators on closed manifolds + theory of Fredholm operators [J,P].

Theorem. Δ is positive. In other words: $(\Delta\omega, \omega) \geq 0$ where $\omega \in \Omega(M)$.

Proof. $(\Delta\omega, \omega) = ((d\delta + \delta d)\omega, \omega) = (d\delta\omega, \omega) + (\delta d\omega, \omega) = (\delta\omega, \delta\omega) + (d\omega, d\omega) = \|\delta\omega\|^2 + \|d\omega\|^2 \geq 0$

Corollary. $(\Delta\omega, \omega) = 0$ iff $\delta\omega = d\omega = 0$, i.e., harmonic forms are both closed and coclosed.

Theorem Poisson's equation $\Delta\psi = \omega_k$ has a solution iff ω_k is orthogonal to $Harm^k(M)$. The solution ψ is noted as $\psi = \Delta^{-1}\omega$.

Proof. The theory of Elliptic operators explains the sufficiency [J]. The necessity is shown as follows:

If there exists ψ such that $\Delta\psi = \omega_k$ then taking γ from $Harm^k(M)$ we have:

$$(\omega, \gamma_k) = (\Delta\psi, \gamma_k) = (\psi, \Delta\gamma_k) = (\psi, 0) = 0$$

which shows that ω_k is orthogonal to $Harm^k(M)$.

Theorem. Harmonic, Exact and co-exact forms are mutually orthogonal spaces.

Proof. Since $d^2 = 0$ then $(d^2\alpha_{k-1}, \beta_{k+1}) = (d\alpha_{k-1}, \delta\beta_{k+1}) = 0$, showing that exact and coexact forms are orthogonal.

Since harmonic forms are closed then if γ_k is harmonic and

$d\gamma_k = 0$ hence $(\beta_{k+1}, d\gamma_k) = (\delta\beta_{k+1}, \gamma_k) = 0$ showing the mutual orthogonality between coexact and harmonic forms.

likewise $\delta\gamma_k = 0$ hence $(\alpha_{k-1}, \delta\gamma_k) = (d\alpha_{k-1}, \gamma_k) = 0$ showing the mutual orthogonality between exact and harmonic forms.

Hodge Decomposition Theorem. Harmonic, Exact and co-exact forms generate $\Omega^k(M)$: $\Omega^k(M) = Ker d \oplus Ker d^* \oplus Ker \Delta$.

Proof. Let $P : \Omega^k(M) \rightarrow \text{Harm}^k(M)$ be the orthogonal projection operator generated by the scalar product of forms, then for any $\omega \in \Omega^k(M)$ we have that $\omega - P\omega$ is orthogonal to $\text{Harm}^k(M)$. Hence, Poisson's equation

$$\Delta\psi = \omega - P\omega \text{ has a solution that can be written as } \psi = \Delta^{-1}(\omega - P\omega)$$

In conclusion, it makes sense to write

$$\omega = \Delta\psi + P\omega = (d\delta + \delta d)\psi + P\omega = d(\delta\psi) + \delta(d\psi) + P\omega$$

that reads: any form can be orthogonally decomposed as a sum of an exact form plus a coexact form plus a harmonic form.

Hodge Theorem. $H^k(M) = \text{Harm}^k(M)$, in other words, any cohomology class is the class of a harmonic form and every harmonic form is a nontrivial member of $H^k(M)$.

Proof. Our universe is the space of closed forms, since $H^k(M)$ is the space of equivalence classes of closed forms defined by the relation $\omega \sim \omega'$ if there exists $d\psi$ such that $\omega = \omega' + d\psi$.

Let us show that there is an isomorphism between $H^k(M)$ and $\text{Harm}^k(M)$ induced by the projection operator over harmonic forms $P : \Omega^k(M) \rightarrow \text{Harm}^k(M)$. The isomorphism is

$$P([\omega]) = [P\omega].$$

Decomposing ω in its components:

$$\omega = d(\delta\psi) + \delta(d\psi) + P\omega$$

but $d\omega = 0$ hence applying d to both sides of this equation

$$d\omega = d^2(\delta\psi) + d\delta(d\psi) + dP\omega = d\delta(d\psi) + dP\omega$$

but $dP\omega = 0$ because $P\omega$ is harmonic and harmonic forms are closed, so

$$d\omega = d\delta(d\psi) = 0$$

$$\text{hence } (d\delta(d\psi), d\psi) = (\delta(d\psi), \delta(d\psi)) = 0 \text{ hence } \delta d\psi = 0.$$

Therefore

$$\omega = d(\delta\psi) + P\omega.$$

Thus, any closed form can be decomposed into an exact and a harmonic form. Moreover, we can clearly specify ψ : since $\omega - P\omega = d(\delta\psi)$ is exact, it is orthogonal to harmonic forms, and so the corresponding Poisson's equation has a solution. This decomposition can be translated into the language of cohomology classes:

The closed form ω is in its class $[\omega]$ and another representative reads $\omega' = \omega + d\phi$. Proceeding as before we find

$$\omega + d\phi = d(\delta\psi') + P(\omega + d\phi)$$

But $d\phi$ is exact, so it has no harmonic component:

$$P(\omega + d\phi) = P\omega + Pd\phi = P(\omega)$$

which means that if two closed forms belong to the same cohomology class, they have the same harmonic component. Therefore, the projector is well defined over cohomology classes.

Let us now prove that different cohomology classes correspond to different harmonic forms.

Let us suppose that $P([\omega]) = P([\eta])$. Let us decompose both forms:

$$\omega = d(\delta\psi) + P\omega$$

$$\eta = d(\delta\rho) + P\eta$$

then

$$\omega - \eta = d(\delta\psi) + P\omega - d(\delta\rho) + P\eta = d(\delta\psi) - d(\delta\rho) = d(\delta\psi - \delta\rho)$$

and we have discovered that if two forms have the same harmonic component, then they are related by an exact form, and so they belong to the same cohomology class. Nice.

Let us show now that any harmonic form is a nontrivial member of $H^k(M)$, that is for $\gamma \in \text{Harm}^k(M)$ we have $d\gamma = 0$ and $\nexists\psi$ such that $\gamma = d\psi$. The first requirement is automatically fulfilled because any harmonic form is closed and the second relies on the fact that harmonic forms and exact forms are orthogonal one to another.

Commutation properties for $*$, d , δ

We can decompose δ as $\delta = - * d *$. Observe that δ takes k -forms and produces $k-1$ -forms. In this respect, it is similar to integration. Indeed, over p -forms $*$ produces an $(n-p)$ -form over which d produces a $(n-p+1)$ -form and hence $\delta = - * d *$ produces a $n - (n-p+1) = p-1$ form. Henceforth, recalling that $** = (-1)^{(n-p+1)(p-1)}$ we have that

$$*\delta = - * d * = (-1)^{(n-p+1)(p-1)+1} d *$$

because $*$ produces and $(n-p)$ -form over which d produces a $(n-p+1)$ -form and hence $- * *$ operates as $(-1)^{(n-p+1)(n-(n-p+1))+1}$. On the other hand, $\delta * = - * d * = - * d (-1)^{p(n-p)} = (-1)^{p(n-p)+1} * d$. Or, $\delta * = (-1)^{p(n-p)+1} * d$. Therefore, multiplying by the sign in the right side we get

$$*d = (-1)^{p(n-p)+1} \delta *.$$

With the help of this identity we can prove the expected result:

Theorem. The wedge product of harmonic forms is harmonic and the Hodge star of a harmonic form is also harmonic.

Proof: Let us take two harmonic forms, γ_1, γ_2 . Both are closed and coclosed, i.e., $d\gamma_1 = d\gamma_2 = \delta\gamma_1\delta\gamma_2 = 0$. Let us calculate the Laplacian of their wedge product:

$$\begin{aligned} \Delta(\gamma_1 \wedge \gamma_2) &= (d\delta + \delta d)(\gamma_1 \wedge \gamma_2) = \\ &= d\delta(\gamma_1 \wedge \gamma_2) + (\delta d)(\gamma_1 \wedge \gamma_2) = \\ &= d(\delta\gamma_1 \wedge \gamma_2 \pm \gamma_1 \wedge \delta\gamma_2) + \delta(d\gamma_1 \wedge \gamma_2 \pm \gamma_1 \wedge d\gamma_2) = 0. \end{aligned}$$

It rests to prove now that if γ is harmonic, so it is $*\gamma$, i.e., if $\Delta(\gamma) = 0$ then $*\Delta(\gamma) = 0$.

$$\begin{aligned} *\Delta(\gamma) &= (\delta d + d\delta)(* \gamma) \\ &= (\delta d * + d\delta *) \gamma = ((\delta(-1)^{(n-p+1)(p-1)+1} + d(*d(-1)^{p(n-p)+1})) \gamma \\ &= (*d\delta(-1)^{(n-p+1)(p-1)+1}(-1)^{(p-1)(n-p+1)+1} + *\delta d(-1)^{p(n-p)+1}(-1)^{(n-p)p+1}) \gamma \\ &= (*(-1)^{2(n-p+1)+2} d\delta + \delta d(-1)^{2p(n-p)+2}) \gamma \\ &= (*(\delta d + d\delta)) \gamma. \end{aligned}$$

We have thus proved

Theorem $\text{Harm}(\Omega)$ yields a space of representatives of $H^*(\Omega)$ that conserves all its algebraic properties.

Proof. The projection is a continuous operator that is also a retraction.

Theorem. Let E be the base of $Harm^{2k}$ formed with those elements of H^{2k} that are harmonic. Then, the signature quadratic form Q is block diagonal in $\{E\}$ and the signature of M fulfills $\sigma(M) = \dim(Harm_+^{2k}) - \dim(Harm_-^{2k})$ where $Harm_+^{2k}, Harm_-^{2k}$ are the eigenspaces of $*$ with ± 1 eigenvalues.

Proof. Over $2k$ -forms in a manifold of dimension $4k$, the Hodge $*$ satisfies $** = 1$. Hence $*$ has eigenvalues ± 1 . Let $*\omega^+ = \omega^+$ and $*\omega^- = -\omega^-$. Then:

$$\begin{aligned}\sigma(\omega^+, \omega^+) &= \int \omega^+ \wedge \omega^+ = \int \omega^+ \wedge *\omega^+ = (\omega^+, \omega^+) \geq 0 \\ \sigma(\omega^-, \omega^-) &= \int \omega^- \wedge \omega^- = - \int \omega^- \wedge *\omega^- = -(\omega^-, \omega^-) \leq 0\end{aligned}$$

Using the fact that $\alpha \wedge *\beta = \beta \wedge *\alpha$ and that in our case the wedge product is commutative we have

$$\sigma(\omega^+, \omega^-) = \int \omega^+ \wedge \omega^- = - \int \omega^+ \wedge *\omega^- = - \int \omega^- \wedge *\omega^+ = - \int \omega^- \wedge \omega^+ = -\sigma(\omega^-, \omega^+)$$

$$\text{So, } \sigma(\omega^+, \omega^-) = -\sigma(\omega^-, \omega^+) = 0.$$

These results show that the matrix of Q evaluated over a base of harmonic forms is block diagonal, say, D . This allows to alternatively define the signature of a manifold using the following idea: for a real number r , we can define its sign as $sign(r) = r/\sqrt{r^2}$. For selfadjoint diagonal matrices we also have $sign(D) = D(\sqrt{D^2})^{-1}$. In a general case $sign(A) = A(\sqrt{A^*A})^{-1}$ which displays a diagonal filled in ± 1 . Thus

$$\sigma(M) = \text{Trace}(sign(Q))$$

where Q is any matrix representing the signature quadratic form.

When some eigenvalues of a matrix are zero, they don't enter to define the signature of a given matrix. So, it would be nice to face up at this point the following intrigue: Is Q degenerate allowing the existence of null eigenvalues?

Theorem. Q is not degenerate over H^* .

Proof. It is enough to take a harmonic representative, γ which is both closed and coclosed with $*\gamma$ also harmonic. Let us suppose now that

$\int(\gamma_1 \wedge \gamma_2) = 0, \forall \gamma_2$. We shall prove that then $\gamma_1 = 0$. If that is true, then we can take $\gamma_1 = *\gamma_2$, therefore we get

$$\int(\gamma_1 \wedge *\gamma_1) = (\gamma_1, *\gamma_1) = 0$$

from which we conclude that $\gamma_1 = 0$, which means that Q is not degenerate and that has no zero eigenvalues.

Moreover, if we note $Harm^\pm$ as the eigenspaces of $*$ with ± 1 eigenvalues, we can conclude that the signature of this matrix is the number of positive eigenvalues minus the number of negative eigenvalues but this subtraction corresponds to

$$\sigma(M) = \dim(Harm^+) - \dim(Harm^-)$$

Challenge. Since the index of a (Fredholm) operator P is $ind(P) = \dim(Ker P) - \dim(P^*)$ where P^* is the adjoint of P , this last equality seems to

predict the existence of an operator (over an infinite dimensional Hilbert space) whose index is precisely the signature of the manifold.

Lemma. Let $\dim M = 4k$ then the dimension of H^{2k} and the signature of M have the same parity, i.e., $b_{2k} = \sigma(M) \bmod 2$.

Since the harmonic forms $Harm^{2k}$ can be decomposed into the direct sum $Harm^{2k} = Harm_+^{2k} \oplus Harm_-^{2k}$ thus $b_{2k} = \dim(Harm_+^{2k}) + \dim(Harm_-^{2k})$ and moreover,

$$\sigma(M) = \dim(Harm_+^{2k}) - \dim(Harm_-^{2k})$$

Therefore

$$b_{2k} = \dim(Harm_+^{2k}) - \dim(Harm_-^{2k}) + 2\dim(Harm_-^{2k}) = \sigma(M) + 2\dim(Harm_-^{2k})$$

$$\text{or } b_{2k} = \sigma(M) + 2\dim(Harm_-^{2k})$$

$$\text{or } b_{2k} - \sigma(M) = 2\dim(Harm_-^{2k})$$

$$\text{or } b_{2k} = \sigma(M) \bmod 2.$$

Since "to have the same parity" is transitive, we can conclude

Corollary. Let $\dim M = 4k$ then the signature of M and the Euler characteristic has the same parity, i.e., $\sigma(M) = \chi(M) \bmod 2$.

Remark. The equalities mod 2 that we have found point to the possibility of relating the signature operator with the Pontrjagin classes in which coefficients are taken from Z_2 .

4 The signature operator

The following properties of the Hodge-star $*$ operator are necessary for our purposes in this section:

1. $** = (-1)^{p(n-p)}$ when $*$ operates over a p -form. Hence, over $(p+1)$ -forms $** = (-1)^{(p+1)(n-p-1)}$ while over $(p-1)$ -forms $** = (-1)^{(p-1)(n-p+1)}$.

2. Let us define the chirality operator $J = i^{n/2+p(p-1)}*$ when it operates over p -forms, which becomes $J = i^{n/2+(p+1)(p)}*$ over $(p+1)$ -forms and $J = i^{n/2+(p-1)(p-2)}*$ over $(p-1)$ -forms. Warning: real scalars are replaced by a complexified version.

3. Recall that operating over p -forms we have

$$*\delta = - * * d * = (-1)^{(n-p+1)(p-1)+1} d *$$

and that

$$*d = (-1)^{p(n-p)+1} \delta *.$$

These identities will be used to change the relative position of $*$ with respect to d, δ .

Theorem. If D is defined as $D = d + \delta$ then J and D anticommute: $JD = -DJ$

Proof: $JD = J(d + \delta) = Jd + J\delta$. Recall that d rises the order of a form by one unit, while δ lowers the order of a form by one. This must be kept in

mind when operating with J : $Jd + J\delta = i^{n/2+(p+1)(p)} * d + i^{n/2+(p-1)(p-2)} * \delta$.

Changing the relative position of $*$ we have:

$$\begin{aligned} JD &= i^{n/2+(p+1)(p)} (-1)^{p(n-p)+1} \delta * + i^{n/2+(p-1)(p-2)} (-1)^{(n-p+1)(p-1)+1} d * \\ &= i^{n/2+(p-1+2)(p)} (-1)^{p(n-p)+1} \delta * + i^{n/2+(p-1)(p)-2(p-1)} (-1)^{(n-p+1)(p-1)+1} d * \\ &= i^{n/2+(p-1)(p)} [i^{2p} (-1)^{p(n-p)+1} \delta + i^{-2(p-1)} (-1)^{(n-p+1)(p-1)+1} d] * \\ &= [i^{2p} (-1)^{p(n-p)+1} \delta + i^{-2(p-1)} (-1)^{(n-p+1)(p-1)+1} d] J \\ &= [(-1)^{p+p(n-p)+1} \delta + (-1)^{-(p-1)+(n-p+1)(p-1)+1} d] J. \end{aligned}$$

Now, let us see that if n is even then the exponents in this last expression are odd. To see this, recall that $p(p-1)$ is always even, that if n is even then so is np and that p^2 always behave like p , i.e., module two we have the following equalities:

$$\begin{aligned} p + p(n-p) + 1 &= p + np - p^2 + 1 = p + 0 - p + 1 = 1(\text{odd}). \\ -(p-1) + (n-p+1)(p-1) + 1 &= (p-1)(-1+n-p+1) + 1 = (p-1)(n-p) + 1 = \\ pn - p^2 - n + p + 1 &= 0 - p - 0 + p + 1 = 1(\text{odd}). \end{aligned}$$

Putting all together:

$$JD = -(\delta + d)J = -DJ, \text{ and so, } J \text{ and } D \text{ anticommute.}$$

Theorem. $J^2 = 1$ if $n = 2k$

Proof. We know that if $J = i^{n/2+p(p-1)} *$ operates over p -forms, it produces $n-p$ -forms, so a new application of J operates over $(n-p)$ -forms and hence J takes the form $J = i^{n/2+(n-p)(n-p-1)} *$. Thus

$$J^2 = i^{n/2+(n-p)(n-p-1)} * i^{n/2+p(p-1)} * = i^{n/2+(n-p)(n-p-1)+n/2+p(p-1)} **$$

Recalling that over p -forms $** = (-1)^{p(n-p)} = (i)^{2p(n-p)}$ we get:

$$\begin{aligned} J^2 &= i^{n/2+(n-p)(n-p-1)+n/2+p(p-1)} (i)^{2p(n-p)} \\ &= i^{n/2+(n-p)(n-p-1)+n/2+p(p-1)+2p(n-p)} = i^{n+(n-p)(n-p-1)+p(p-1)+2p(n-p)} \\ &= i^{n+(n-p)(n-p-1)+p(p-1)+2p(n-p)} = i^{n+n^2-np-n-np+p^2+p+p^2-p+2np-2p^2} \end{aligned}$$

Simplifying we obtain

$$J^2 = i^{n^2} = i^{4k^2} = (i^4)^{k^2} = 1.$$

Corollary The eigenvalues of J are ± 1 and for any ω , $J\omega = \pm\omega$.

If $J\omega = \lambda\omega$ then $J^2\omega = \lambda J\omega = \lambda^2\omega = \omega$ then $\lambda^2 = 1$ and $\lambda = \pm 1$.

Lema. D interchanges the eigenspaces of J . In other words, if $J\omega = \omega$ then $JD\omega = -D\omega$ and viceversa if $J\omega = -\omega$ then $JD\omega = D\omega$.

Proof: If $J\omega = \omega$ then $DJ\omega = D\omega$, but $DJ\omega = -JD\omega$ because these operators anticommute, hence $-JD\omega = D\omega$ or $JD\omega = -D\omega$. Likewise, if $J\omega = -\omega$ then $DJ\omega = -JD\omega = -D\omega$ or $JD\omega = D\omega$.

Definition. Let E be the positive sector of J and F the negative sector. The signature operator D^+ of D is the restriction of D to E which goes onto F .

Theorem. The index of the signature operator is equal to the signature of M .

Proof: Since $D = d + \delta$ interchanges eigenspaces of J , we can write

$$D = \begin{pmatrix} 0 & D_- \\ D_+ & 0 \end{pmatrix}$$

where $D_- = D_+^*$. Indeed,
 $(D_+\alpha, \beta) = ((d+\delta)\alpha, \beta) = (\alpha, (d+\delta)^*\beta) = (\alpha, (d^*+\delta^*)\beta) = (\alpha, (\delta+d)\beta) =$
 $(\alpha, D_-\beta)$
 where we have taken care of the domains of definitions. Hence

$$\Delta = D^2 = \begin{pmatrix} D_+^*D_+ & 0 \\ 0 & D_+D_+^* \end{pmatrix} = \begin{pmatrix} \Delta_+ & 0 \\ 0 & \Delta_- \end{pmatrix}$$

where $\Delta_{\pm}$ are the restrictions to the $\pm$ eigenspaces of J of the Laplace operator Δ .

Taking $2k$ -forms and the corresponding operators we have:

$$\text{Ind}(D_+) = \dim(\text{Ker}D_+) - \dim(\text{Ker}D_+^*)$$

Recalling that $\text{ind}(F) = \text{ind}(F^*)$, we can add to the right side

$$0 = \dim(\text{Ker}D_+^*) - \dim(\text{Ker}D_+^*) = \dim(\text{Ker}D_+^*) - \dim(\text{Ker}D_+)$$

and we have

$$\text{Ind}(D_+) = \dim(\text{Ker}D_+) - \dim(\text{Ker}D_+^*) + \dim(\text{Ker}D_+^*) - \dim(\text{Ker}D_+)$$

$$= \dim(\text{Ker}D_+^*) + \dim(\text{Ker}D_+) - (\dim(\text{Ker}D_+) + \dim(\text{Ker}D_+^*))$$

but $\dim(\text{Ker}AB) = \dim(\text{Ker}A) + \dim(\text{Ker}B)$, hence

$$\text{Ind}(D_+) = \dim(\text{Ker}D_+^*D_+) - \dim(\text{Ker}D_+D_+^*)$$

$$= \dim(\text{Ker}\Delta_+) - \dim(\Delta_-)$$

$$= \dim(\text{Harm}_+^{2k}) - \dim(\text{Harm}_-^{2k}) = \sigma(M)$$

5 References

1. Frankel, Theodore, (1997), The geometry of physics : an introduction. Cambridge, England ; New York : Cambridge University Press, c1997.
2. Joshi M.S., (1999), 'Introduction to Pseudo-differential Operators', arXiv:math.AP/9906155.
3. Nakahara Mikio, (1990). *Geometry, topology, and physics*. Bristol, England ; Institute of Physics Publishing.
4. Paycha S., 1997, "Une petite introduction aux FIBRES DETERMINANTS Pour grands débutants". Apuntes Matemáticos, No. 38, Departamento de Matemáticas, Uniandes., Bogotá, Colombia.

6 Acknowledgments

Sylvie Paycha